

# Chapter 111

## Beyond a Simple Effect: Variable and Changing Responses to Anthropogenic Noise

Andrew N. Radford, Julia Purser, Rick Bruintjes, Irene K. Voellmy, Kirsty A. Everley, Matthew A. Wale, Sophie Holles, and Stephen D. Simpson

**Abstract** A growing number of experimental studies have demonstrated that exposure to anthropogenic noise can affect the behavior and physiology of a variety of aquatic organisms. However, work in other fields suggests that responses are likely to differ between species, individuals, and situations and across time. We suggest that issues such as interspecific and intrapopulation variation, context dependency, repeated exposure and prior experience, and recovery and compensation need to be considered if we are to gain a full understanding of the impacts of this global pollutant.

**Keywords** Condition dependency • Context dependency • Interspecific variation • Repeated exposure • Recovery

### 1 Introduction

Human activities such as urban development, the construction and exploitation of natural resources, and transportation have increased around the globe in the last century, changing the acoustic environment both on land and underwater (Jasny 1999; McDonald et al. 2006; Watts et al. 2007; Barber et al. 2009). In addition to the unprecedented modification of the natural soundscape, the nature of the sound generated by human activities is often very different from that arising from natural sources; anthropogenic noises may differ from abiotic or biotic sounds in such acoustic characteristics as constancy, rise time, duty cycle, and

---

A.N. Radford (✉) • J. Purser • R. Bruintjes • I.K. Voellmy • M.A. Wale • S. Holles  
School of Biological Sciences, University of Bristol, Woodland Road, Bristol BS8 1UG, UK  
e-mail: [andy.radford@bristol.ac.uk](mailto:andy.radford@bristol.ac.uk); [julia.purser@bristol.ac.uk](mailto:julia.purser@bristol.ac.uk); [rick.bruintjes@bristol.ac.uk](mailto:rick.bruintjes@bristol.ac.uk);  
[irene.vollmy@bristol.ac.uk](mailto:irene.vollmy@bristol.ac.uk); [matt.wale@bristol.ac.uk](mailto:matt.wale@bristol.ac.uk); [sophie.holles@bristol.ac.uk](mailto:sophie.holles@bristol.ac.uk)

K.A. Everley • S.D. Simpson  
Department of Biosciences, College of Life and Environmental Sciences,  
University of Exeter, Exeter EX4 4QD, UK  
e-mail: [s.simpson@exeter.ac.uk](mailto:s.simpson@exeter.ac.uk)

impulsiveness (Hildebrand 2009; Popper and Hastings 2009). Consequently, anthropogenic noise presents a very real, and often novel, challenge to animals and is now recognized as a pollutant of international concern (e.g., inclusion in the US National Environment Policy Act and the European Commission Marine Strategy Framework Directive and as a permanent item on the agenda of the International Maritime Organization's Marine Environmental Protection Committee).

In recent years, anthropogenic noise has been demonstrated to affect a variety of taxonomic groups across a range of scales, from the physiology and behavior of individuals to changes at the population and community level (see Tyack 2008; Barber et al. 2009; Popper and Hastings 2009; Slabbekoorn et al. 2010; Kight and Swaddle 2011 for reviews). Experimental studies on aquatic organisms have indicated that mammals, fishes, and invertebrates are all vulnerable. For instance, noise causes alterations in the vocal signaling, movement patterns, foraging behavior, and hearing thresholds of marine mammals (e.g., Bejder et al. 2006; Mooney et al. 2009; Di Iorio and Clark 2010; Tyack et al. 2011). In fishes, movement, settlement, foraging, social interactions, and antipredator behavior are all influenced by anthropogenic noise (e.g., Purser and Radford 2011; Bruintjes and Radford 2013; Holles et al. 2013; see Chapters 32 by Everley et al. and 129 by Simpson et al.), which has also been shown to cause stress, temporary threshold shifts, and injury (e.g., Smith et al. 2004; Wysocki et al. 2006; Halvorsen et al. 2012). Marine invertebrates, which use sound for a variety of reasons (e.g., Simpson et al. 2011; Vermeij et al. 2010), are also impacted because anthropogenic noise has both physiological and behavioral consequences (e.g., Wale et al. 2013a, b).

## 2 Beyond a Simple Effect

Understandably for an emerging research field (the majority of studies have been published in the last 3 years; reviewed in Radford et al. 2012; Morley et al. 2014), the main question considered in experimental studies to date has been the straightforward, yet important, "Does anthropogenic noise have an impact?" We argue that attention also needs to focus on additional questions relating to variable and changing responses to noise, and we outline some examples below.

### 2.1 *Interspecific Differences*

It is likely that there will be stable interspecific differences in susceptibility and responses to elevated noise levels depending on variation in, for example, hearing ability (Fay et al. 2008) and mechanisms of physiological stress response (Hofer and East 1998). Direct comparisons of species in response to the same noise

source in the same contexts are rare. However, Halvorsen et al. (2012) recently showed variation in the level of injury caused in different fish species by exposure to the same pile-driving stimulus. Voellmy (2013) has also demonstrated that the foraging and antipredator behavior of three-spined sticklebacks (*Gasterosteus aculeatus*) and European minnows (*Phoxinus phoxinus*) is affected differently by the same noise playbacks. In sympatry, and particularly if there is an overlap in ecological niches, such differences may affect the relative success of each species under scenarios of disturbance and so potentially may affect community composition and structure. These effects could arise through alterations in the interactions between, for example, competitors, predators and prey, and plants and pollinators (see Francis et al. 2009, 2012).

## 2.2 *Intrapopulation Variation*

It is clear from other research fields that factors such as sex, dominance status, age, size, and condition may all influence how members of the same population are affected by a given stimulus, including environmental change arising from human activities (Kiffney and Clements 1996; Huntingford et al. 2006; Xu et al. 2010). Although empirical work on the impacts of anthropogenic noise has tended to focus on the overall response of cohorts of individuals, studies are beginning to consider potential intrapopulation differences in response. For instance, Wale et al. (2013a) found that heavier shore crabs (*Carcinus maenas*) consume a disproportionately larger amount of oxygen than smaller individuals when exposed to playback of ship noise. Brintjes and Radford (2013) discovered that dominant male and female individuals of the cooperatively breeding cichlid fish species *Neolamprologus pulcher* exhibit different behavioral responses to the same playback of boat noise. More recently, Purser et al. (in preparation) have shown that European eels (*Anguilla anguilla*) in poorer body condition (relative weight) suffer more detrimental physiological and behavioral consequences than individuals in better condition. Consistent interindividual differences in response could have impacts on population dynamics and for harvests of commercially important species.

## 2.3 *Context-Dependent Responses*

The response of an animal can be dependent on its current situation (e.g., Bell et al. 2009, 2010), with increasing evidence that context can influence the harmful effects of human activities on animal welfare (see Huntingford et al. 2006 and references therein). Recent work by Brintjes and Radford (2013) showcases that the impact of anthropogenic noise can be context dependent; playback of boat noise resulted in a reduction in antipredator defense by *Neolamprologus pulcher* group members if no

eggs were present in a nest but not if eggs were present. Moreover, social interactions between dominants and subordinates were affected differently by the same noise playbacks depending on whether group members were engaged in defense behavior or nest digging. The implication is that responses to anthropogenic noise are not fixed but rather show some element of flexibility, which may or may not be under the control of the individual.

## ***2.4 Repeated Exposure and Prior Experience***

Responses to pollutants may change across time (Piola and Johnston 2009; Whitehead et al. 2010) as a result of such processes as habituation, tolerance, and sensitization (Bejder et al. 2009). Although noise-related experiments have, from an understandable logistical perspective, often involved a single presentation of the relevant stimulus, organisms in most natural situations are likely to experience chronic or repeated noise exposure. Exploring how responses can change and are dependent on prior experience is thus important (Simpson et al. 2010; see Chapter 149 by Voellmy et al.). Wale et al. (2013a) showed that although shore crabs repeatedly exposed to ambient-noise playback increased their oxygen consumption (perhaps due to handling stress), those individuals repeatedly exposed to playback of ship noise did not exhibit a similar change. It is possible that they had already shown a maximum response on a first exposure to ship-noise playback, but they might also have become habituated or tolerant over time. Voellmy (2013) has also recently demonstrated, by manipulating holding-tank noise conditions, that prior acoustic experience can influence responses of fish to experimental playbacks, whereas Nedelec et al. (2014; in preparation) have found that repeated exposure to boat-noise playback can affect subsequent behavioral and physiological responses in the early life stages of both fish and marine invertebrates.

## ***2.5 Recovery and Compensation***

Many anthropogenic noise events are transient in nature (Hildebrand 2009; Popper and Hastings 2009), and short-term impacts of noise may not necessarily translate into long-term consequences (see Bejder et al. 2006). Although it is clear that behavior and physiology can be detrimentally impacted during the period of elevated noise (see Section 1), the effects on survival and reproductive success will be dependent on whether, and how quickly, the affected individuals recover to baseline performance levels and if they can compensate. Bruintjes et al. (in preparation) have recently found that the detrimental effects of ship-noise playback on European eel antipredator behavior and respiratory rate are not sustained when the noise ceases. Species will differ in their ability to recover and compensate (Voellmy 2013), and compensation itself may carry a variety of inherent costs (see Purser and

Radford 2011). Thus, studies are needed that look at longer time frames and consider postexposure periods as well as when the noise itself is apparent.

### 3 Conclusions

The human population is projected to increase by 2.3 billion between 2011 and 2050 (United Nations 2011) and thus noise pollution is not just a pressing issue but one of ever-increasing concern. It is now well established that anthropogenic noise does indeed impact a wide range of animals. As the research field moves forward, we advocate the exploration of a wider range of questions such that we can understand more fully the range of effects of this global pollutant and thus optimize strategies to mitigate impacts to both aquatic and terrestrial ecosystems.

**Acknowledgments** We are grateful to the Department for Environment, Food, & Rural Affairs (Defra), the Natural Environment Research Council (NERC), the Engineering and Physical Sciences Research Council (EPSRC), the Swiss National Science Foundation, the Association for the Study of Animal Behaviour (ASAB), Marine Scotland, Subacoustech, the Professional Association of Diving Instructors (PADI) Aware, and the University of Bristol for funding our research and facilitating our discussions.

### References

- Barber JR, Crooks KR, Fristrup KM (2009) The costs of chronic noise exposure for terrestrial organisms. *Trends Ecol Evol* 25:180–189
- Bejder L, Samuels A, Whitehead H, Finn H, Allen S (2009) Impact assessment research: use and misuse of habituation, sensitisation and tolerance in describing wildlife responses to anthropogenic stimuli. *Mar Ecol Prog Ser* 395:177–185
- Bejder L, Samuels A, Whitehead H, Gales N (2006) Interpreting short-term behavioural responses to disturbance within a longitudinal perspective. *Anim Behav* 72:1149–1158
- Bell MBV, Radford AN, Rose R, Wade HM, Ridley AR (2009) The value of constant surveillance in a risky environment. *Proc R Soc Lond B Biol Sci* 276:2997–3005
- Bell MBV, Radford AN, Smith RA, Thompson AM, Ridley AR (2010) Bargaining babblers: vocal negotiation of cooperative behavior in a social bird. *Proc R Soc Lond B Biol Sci* 277:3223–3228
- Bruintjes R, Radford AN (2013) Context-dependent impacts of anthropogenic noise on individual and social behavior in a cooperatively breeding fish. *Anim Behav* 85:1343–1349
- Di Iorio L, Clark CW (2010) Exposure to seismic survey alters blue whale acoustic communication. *Biol Lett* 6:51–54
- Fay RR, Popper AN, Webb JF (2008) Introduction to fish bioacoustics. In: Webb JF, Fay RR, Popper AN (eds) *Fish bioacoustics*. Springer, New York, pp 1–15
- Francis CD, Kleist NJ, Ortega CP, Cruz A (2012) Noise pollution alters ecological services: enhanced pollination and disrupted seed dispersal. *Proc R Soc Lond B Biol Sci* 279:2727–2735
- Francis CD, Ortega CP, Cruz A (2009) Noise pollution changes avian communities and species interactions. *Curr Biol* 19:1415–1419

- Halvorsen MB, Casper BM, Matthews F, Carlson TJ, Popper AN (2012) Effects of exposure to pile-driving sounds on the lake sturgeon, Nile tilapia and hogchoker. *Proc R Soc Lond B Biol Sci* 279:4705–4714
- Hildebrand JA (2009) Anthropogenic and natural sources of ambient noise in the ocean. *Mar Ecol Prog Ser* 395:4–20
- Hofer H, East ML (1998) Biological conservation and stress. *Adv Study Behav* 27:405–525
- Holles S, Simpson SD, Radford AN, Berten L, Lecchini D (2013) Boat noise disrupts orientation behaviour in a coral reef fish. *Mar Ecol Prog Ser* 485:295–300
- Huntingford FA, Adams C, Braithwaite VA, Kadri S, Pottinger TG, Sandoe P, Turnbull JF (2006) Current issues in fish welfare. *J Fish Biol* 68:332–372
- Jasny M (1999) Sounding the depths: supertankers, sonar, and the rise of undersea noise. Natural Resources Defense Council, New York
- Kiffney P, Clements W (1996) Size dependent response of macroinvertebrates to metals in experimental streams. *Env Toxicol Chem* 15:1352–1356
- Kight CR, Swaddle JP (2011) How and why environmental noise impacts animals: an integrative, mechanistic review. *Ecol Lett* 14:1052–1061
- McDonald MA, Hildebrand JA, Wiggins SM (2006) Increases in deep ocean ambient noise in the Northeast Pacific west of San Nicholas Island, California. *J Acoust Soc Am* 120:711–718
- Mooney TA, Machtigall PE, Vlachos S (2009) Sonar-induced temporary hearing loss in dolphins. *Biol Lett* 5:565–567
- Morley EL, Jones G, Radford AN (2014) The importance of invertebrates when considering the impacts of anthropogenic noise. *Proc R Soc Lond B Biol Sci* 281:20132683, <http://dx.doi.org/10.1098/rspb.2013.2683>
- Nedelec SL, Radford AN, Simpson SD, Nedelec B, Lecchini D, Mills SC (2014) Anthropogenic noise playback impairs embryonic development and increases mortality in a marine invertebrate. *Sci Rep* 4:5891
- Piola RF, Johnston EL (2009) Comparing differential tolerance of native and non-indigenous marine species to metal pollution using novel assay techniques. *Environ Pollut* 157:2853–2864
- Popper AN, Hastings MC (2009) The effects of anthropogenic sources of sound on fishes. *J Fish Biol* 75:455–489
- Purser J, Radford AN (2011) Acoustic noise induces attention shifts and reduces foraging performance in three-spined sticklebacks (*Gasterosteus aculeatus*). *PLoS ONE* 6:e17478
- Radford AN, Morley EL, Jones G (2012) The effects of noise on biodiversity, Defra Report NO0235. Department for Environment, Food, & Rural Affairs, London
- Simpson SD, Meekan MG, Larsen NJ, McCauley RD, Jeffs A (2010) Behavioural plasticity in larval reef fish: orientation is influenced by recent acoustic experiences. *Behav Ecol* 21:1098–1105
- Simpson SD, Radford AN, Tickle EJ, Meekan MG, Jeffs AG (2011) Adaptive avoidance of reef noise. *PLoS ONE* 6:e16625
- Slabbekoorn H, Bouton N, van Opzeeland I, Coers A, ten Cate C, Popper AN (2010) A noisy spring: the impact of globally rising underwater sound levels on fish. *Trends Ecol Evol* 25:419–427
- Smith ME, Kane AS, Popper AN (2004) Noise-induced stress response and hearing loss in goldfish (*Carassius auratus*). *J Exp Biol* 207:427–435
- Tyack P (2008) Implications for marine mammals of large-scale changes in the marine acoustic environment. *J Mamm* 89:549–558
- Tyack PL, Zimmer WMX, Moretti D, Southall BL, Claridge DE, Durban JW, Clark CW, D'Amico A, DiMarzio N, Jarvis S, McCarthy E, Morrissey R, Ward J, Boyd IL (2011) Beaked whales respond to simulated and actual naval sonar. *PLoS ONE* 6:e17009
- United Nations (2011) World urbanization prospects: the 2011 revision. Population Division, Department of Economic and Social Affairs, United Nations, New York, [http://esa.un.org/unup/pdf/WUP2011\\_Highlights.pdf](http://esa.un.org/unup/pdf/WUP2011_Highlights.pdf)
- Vermeij MJA, Marhaver KL, Huijbers CM, Nagelkerken I, Simpson SD (2010) Coral larvae move toward reef sounds. *PLoS ONE* 5:e10660

- Voellmy I (2013) Effects of anthropogenic noise on fish behaviour. PhD thesis, University of Bristol, Bristol
- Wale MA, Simpson SD, Radford AN (2013a) Size-dependent physiological responses of shore crabs to single and repeated playback of ship noise. *Biol Lett* 9:20121103
- Wale MA, Simpson SD, Radford AN (2013b) Noise negatively affects foraging and antipredator behaviour in shore crabs. *Anim Behav* 86:111–118
- Watts RD, Compton RW, McCammon JH, Rich CL, Right SM, Owens T, Ouren DDS (2007) Roadless space of the conterminous United States. *Science* 316:736–738
- Whitehead A, Triant DA, Champlin D, Nacci D (2010) Comparative transcriptomics implicates mechanisms of evolved pollution tolerance in a killifish population. *Mol Ecol* 19:5186–5203
- Wysocki LE, Dittami JP, Ladich F (2006) Ship noise and cortisol secretion in European freshwater fishes. *Biol Conserv* 128:501–508
- Xu CL, Letcher BH, Nislow KH (2010) Size-dependent survival of brook trout *Salvelinus fontinalis* in summer: effects of water temperature and stream flow. *J Fish Biol* 76:2342–2369